

Observational evidence for summer rainfall at Titan's north pole

Rajani D. Dhingra¹, Jason W. Barnes¹, Robert H. Brown², Bonnie J.

Buratti³, Christophe Sotin³, Philip D. Nicholson⁴, Kevin H. Baines⁵, Roger

N. Clark⁶, Jason M. Soderblom⁷, Ralf Jaumann⁸, Sebastien Rodriguez⁹,

Stéphane Le Mouélic¹⁰, Elizabeth P. Turtle¹¹, Jason E. Perry¹², Valeria

Cottini¹³, and Don E. Jennings¹⁴

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1029/2018GL080943

Rajani Dhingra, rhapsodyraj@gmail.com, rdhingra@uidaho.edu

¹Department of Physics, University of
Idaho, 875 Perimeter Drive, Moscow, ID
83843, USA

² Dept. of Planetary Sciences, University
of Arizona, AZ, USA

³Jet Propulsion Laboratory, Caltech, CA,
USA

⁴Cornell University, Astronomy Dept.,
NY, USA

⁵Space Science & Engineering Center,
University of Wisconsin-Madison, WI, USA

⁶U.S.G.S., Denver, CO, USA

⁷Dept. of Earth, Atmospheric and
Planetary Sciences, MIT, MA, USA

⁸Deutsches Zentrum für Luft- und
Raumfahrt, 12489, Germany

⁹Institut de Physique du Globe de Paris
(IPGP)

Methane rain on Saturn's moon Titan makes it the only place, other than Earth, where rain interacts with the surface. When and where that rain wets the surface changes seasonally in ways that remain poorly understood. Here, we report the discovery of a bright ephemeral feature covering an area of 120,000 km² near Titan's north pole in observations from *Cassini's* near-infrared instrument, VIMS on June 7, 2016. Based on the overall brightness, spectral characteristics, and geologic context, we attribute this new feature to specular reflections from a rain-wetted solid surface like those off of a sunlit wet sidewalk. The reported observation is the first documented rainfall event at Titan's north pole and heralds the arrival of the northern summer (through climatic evidence), which has been delayed relative to model predictions. This

¹⁰Laboratoire de Planetologie et
Geodynamique, CNRS UMR6112,
Universite de Nantes, France

¹¹John Hopkins Applied Physics
Laboratory, MD, USA

¹²University of Arizona, AZ, USA

¹³John Hopkins University, MD, USA

¹⁴ Goddard Space Flight Center, MD,
USA

detection helps constrain Titan's seasonal change and shows that the "wet-sidewalk effect can be used to identify other rain events.

Keypoints:

- We report the discovery of a bright ephemeral feature near Titan's north pole in observations from *Cassini's* near-infrared instrument, VIMS.
- We detect a super-bright, broad specular reflection (glint) when the surface is illuminated by the Sun, provided the ground is wet.
- Our detection of a wet surface near Titan's north pole is the first evidence of rain during north polar summer.
- Detections of broad specular reflections is a new technique for monitoring the occurrence of precipitation across Titan's surface.

1. Introduction

A major objective of the *Cassini* Solstice Mission was to search for evidence of seasonal changes on Titan [Tokano *et al.*, 2006; Ádámkovics *et al.*, 2007]—both atmospheric (clouds) and surficial (rainfall)—that were expected to occur as Titan moved toward the northern summer solstice in 2017. Cassini arrived in the Saturn system during southern summer (2004) and observed clouds and rainfall at the south pole [Porco *et al.*, 2005; Schaller *et al.*, 2006; Turtle *et al.*, 2009], consistent with General Circulation Models (GCMs) [Tokano and Lorenz, 2006; Schneider *et al.*, 2012]. However, as northern summer approached, the increase in cloud activity in the northern hemisphere that was predicted by such models was not observed. In particular, models predicted an increase in the cloud, storm, and rain activity by 2016. Instead, only small, isolated patches of clouds over the north pole began appearing in 2014 (T104 and later flybys).

While both *Cassini* and Earth-based telescopes [Schaller *et al.*, 2006; Roe *et al.*, 2002] have observed clouds on Titan, rain has only been inferred through observed surface changes that require high resolution views. Although four instruments on *Cassini* were able to observe Titan's surface: the RADAR [Elachi *et al.*, 2004], Imaging Science Subsystem (ISS) [Porco *et al.*, 2004], Visual and Infrared Mapping Spectrometer (VIMS) [Brown *et al.*, 2004], and Composite Infrared Spectrometer (CIRS) [Jennings *et al.*, 2017], only the VIMS and ISS instruments operate at short enough wavelengths to observe surficial veneers like rainfall.

The ISS instrument observed Titan's surface through the CB3 filter at $0.93\ \mu\text{m}$, and detected surface darkenings associated with rainfall events at southern high latitudes

near the beginning of the mission (southern summer) [Turtle *et al.*, 2009] and later at equatorial latitudes close to the spring equinox [Turtle *et al.*, 2011]. Subsequent to each of these events, VIMS observed surface brightening over months to years after the presumed rainfall [Barnes *et al.*, 2013b]. In neither case could VIMS detect the initial darkening of the surface (likely due to timing and flyby geometries). CIRS in the past has detected seasonal changes in surface temperature variations [Cottini *et al.*, 2012].

2. Observation

Herein, we report the discovery of an extensive ephemeral feature (Figure 1) near Titan's north pole that was observed by *Cassini* VIMS during the T120 (June 07, 2016) flyby and which had disappeared by the next flyby, T121 (July 25, 2016, with similar observation geometry); 3 Titan days (48 Earth days) later. The ephemeral feature is distinct due to its brightness, especially at longer wavelengths ($5\ \mu\text{m}$) and covers an area of $\sim 120,000\ \text{km}^2$ (spreading across 12 VIMS pixels; bigger than any of the Great Lakes on Earth). The spatial sampling of these pixels is on average $\sim 130\ \text{km}$ in the north-south direction and $\sim 50\ \text{km}$ in the east-west direction. In addition to the spectral data from VIMS, we use data from the RADAR, ISS, and CIRS instruments to constrain the origin and nature of this anomalously bright, ephemeral feature that we henceforth call the "Bright Ephemeral Feature" (BEF). The ISS and VIMS data provide spectral information, RADAR data provide geologic context, while the CIRS data provide insight into the regional temperatures.

The previous flyby, T119 (06 May, 2016), occurred 2 Titan days before the T120 flyby, and showed cloud cover over the north pole in VIMS data (see SOM for an image of T119).

Prior to this, the region had most recently been observed by VIMS during the T106 flyby \sim two years earlier (24 Oct, 2014), and the BEF was not observed (Figure 1A) (See SOM for observation geometries). The ephemeral nature of the feature suggests that the BEF was formed by some transient, short-lived process at the north polar region of Titan, which distinctly brightened the region during T120 flyby. Later, the region resumed its usual spectral characteristics by the next flyby, T121 (Figure 1C). We propose that the BEF represents a broad specular reflection from a wetted surface brought about by rainfall.

3. Method

To study how quickly the Bright Ephemeral Feature (BEF) changed, we compared the region corresponding to the BEF in the T120 flyby with the next flyby, T121, and the prior flyby in which the region was visible, T106. Table 1 shows the observation geometries of the VIMS data cubes used. To rule out the possibility that we did not observe the BEF in T106 and T121 due to unfavorable geometry, we list the distance of the T120 BEF from the specular point in each flyby in Table 1. The distance of BEF from the inferred specular point for T106 and T121 (847 km and 470 km) is marginally different from T120's (632 km). Given the small change in the distance between the BEF and T106, T121 specular points, we would have observed the feature if BEF was present on T106 and T121 flybys. This indicates that we do not see the feature in T106 and T121 flyby due to its ephemerality rather than observation geometry.

We create a VIMS color composite of the observation and overlay it on the RADAR map of the north pole of Titan to infer the geographic location of the BEF in Figure 2.

We then create spectra of different spectroscopic units and compare their brightnesses in Figure 3.

When we compare the spectra of different units from the T120 observation in Figure 3B, namely, BEF, specular, clouds, evaporite, and land, we use spectra from the same cube to minimize the atmospheric effects due to observation geometry (See SOM).

The areas where we see the BEF in VIMS are observed at high phase and emission angles by ISS (at $0.93 \mu\text{m}$) which makes it difficult for ISS to detect any darkening or brightening associated with the region. CIRS observations of the region, before and during/after the rainfall event, do not provide clear indications of a temperature drop due to any rainfall-derived cooling of the region.

4. Hypotheses

Several mechanisms could potentially produce the reported spectral signature in VIMS data: clouds, fog, evaporites, direct specular reflections (i.e., reflections off large smooth surfaces), and broad specular reflections. In the following text, we consider the merits of each of these possibilities and demonstrate how all but the broad specular reflections can be ruled out.

The T120 flyby contains extensive clouds in addition to the BEF that serve as the basis for our spectral comparison. Although VIMS observes from 0.3 to $5.1 \mu\text{m}$ in 352 spectral channels, Titan's surface is only visible through the atmosphere in seven narrow spectral windows, at 0.92 , 1.06 , 1.26 , 1.57 , 2.0 , 2.7 and $5 \mu\text{m}$ [Brown *et al.*, 2004]. VIMS data are expressed in terms of surface reflectance as I/F (where I is the observed radiance and πF is the incident solar irradiance). The surface and clouds on Titan show distinct

spectral responses [Rodriguez et al., 2006, 2009]. We incorporate these spectral response characteristics into a color composite (RGB= 5 μm , 2 μm , and 2.7 μm) [Barnes et al., 2007; Sotin et al., 2012] to facilitate the identification of various spectral units in the observation. Though also bright at 5 μm , clouds are brighter at 2.7 μm [McCord et al., 2006; Rodriguez et al., 2011; Griffith et al., 2009] owing to their high altitude. The wings of the 2 μm band of clouds also indicate altitude. As a result, the color composite effectively distinguishes clouds (purple in Figure 1D) from BEF (orange in Figure 1D).

Low-lying clouds or fog [Brown et al., 2009], however, cannot be entirely ruled out simply based on spectral comparison of the BEF with typical high-altitude clouds. Near-surface fog is difficult to distinguish from bright surface features in spectra without a reliable atmospheric radiative transfer correction. However, both the bright and red (5 μm) spectrum of the BEF (in Titan's atmospheric windows) and the fact that the BEF does not extend over nearby large lakes (where one might expect the humidity to be higher) argues against the possibility of a low-lying fog. Even if the latter case is true, the occurrence of a fog layer necessitates a wet (humid) source underneath. Thus, the wetting of land surface by rain is strongly supported, especially considering the large areal spread of the BEF.

Evaporites, direct specular reflections, and broad specular reflections are all surface phenomena. While broad specular reflections from a liquid surface can occur if the surface is roughened by waves [Barnes et al., 2014] or nitrogen bubbles [Malaska et al., 2017; Hofgartner et al., 2014], a broad specular return from a solid surface requires smoothing of the surface at length scales comparable to the wavelength of light, as would occur if the

surface were wet (like a wet-sidewalk after rainfall). RADAR and ISS data observations (Figure 2) show that the BEF occurs over a solid land surface, bordering known lakes in some places: the north- western edge of the BEF coincides with Lagoda Lacus (equivalent radius of ~ 54 km). However, waves on Lagoda Lacus would be discrete, smaller than the lake area, and cannot cause the bright pixels. The BEF is ~ 3 orders of magnitude larger than the lake.

A commonly observed unit, particularly in the north, are $5 \mu\text{m}$ bright deposits observed to abut many filled and empty lakes that are believed to be evaporitic material, comprising deposits left behind when lakes evaporate (Figure 3B) [Barnes *et al.*, 2011; MacKenzie *et al.*, 2014]. Two lines of logic, however, strongly disfavor evaporite deposits for the BEF. The first is the spatial extent. Evaporites typically only border the edges of lake —the deposit observed along the southern end of Ladoga Lacus spans an area of 300 km^2 (as observed during the T76 flyby, May 08, 2011) compared to the BEF, spanning an area of $\sim 120,000 \text{ km}^2$. Second, and most importantly, no evaporites have been observed to form and then disappear within a window of only a few Titan days, as was observed for the BEF (Fig. 1A, B, C).

Direct specular reflections, which we define as the reflection of light off a surface that is smooth at scales both comparable to the wavelength of light and to the projected image of the Sun on the surface [Stephan *et al.*, 2010; Barnes *et al.*, 2013a; Soderblom *et al.*, 2012], depend on viewing-geometry and result in an unusually high I/F. In the T120 flyby we observe the direct specular reflection from the Sun (I/F of ~ 2 in the $5 \mu\text{m}$ spectral window) from the surface of Xolotlan Lacus at 82.7°N , 78.6°W .

We observe the direct specular reflection of the Sun (I/F of ~ 2 in the $5 \mu\text{m}$ spectral window) in the T120 flyby, at 82.7°N , 78.6°W from the surface of the lake, Xolotlan Lacus (Figure 2C), ~ 632 km away from the BEF. As a sidenote, the specular point seems close to the BEF or wetside walk in the Earth analog (Figure 1E). This is due to the smaller angular distance between the specular point and the BEF as compared to T120 observation.

Even though the BEF does not occur at the specular point, it does show spectral similarities to direct specular reflections on Titan. We compare the spectrum of the BEF with those of sea, clouds, land, evaporites, and the direct specular reflection (Figure 3 A, B), all taken from the same flyby (T120) to minimize atmospheric effects (See Method for observation geometry). The direct specular reflection is the brightest “feature” at $5 \mu\text{m}$ in the observation. The next brightest feature at $5 \mu\text{m}$ (by 50%) is the BEF. A rough surface cannot, on its own, reflect specularly. However, surface wetting coats the irregular surfaces of the rough region resulting in a smoother surface than what is underneath. The spectral brightness and observation geometry immediately indicate that the BEF is a broad specular reflection due to smoothing of a surface that was otherwise dry.

5. Modeling surface roughness

In order to test our hypothesis, we model the BEF’s surface roughness to characterize specular reflections from a macroscopically rough surface (Figure 4). We use a numerical planetary specular model [*Barnes et al.*, 2014; *Soderblom et al.*, 2012] with Gaussian-distributed slopes and azimuthal symmetry, for the same observation geometry as the T120 observation (Figure 4B).

For small roughnesses with RMS slopes of $\sigma \sim 2^\circ$ (Figure 4C), only the direct specular reflection at the specular point appears. As we increase the roughness parameter — σ —the specular reflection spreads over a larger number of pixels, each pixel receiving lesser solar flux, resulting in a fuzzier reflection distributed over a broader region. Over an area equivalent to the BEF, the model-determined σ values indicate that the required slopes to get signals comparable to our observation are $\sim 12 - 14^\circ$ (Figure 4D). This is twice the σ value of the inferred waves in Punga Mare (liquid surface), which were determined using the same model, to be $\sim 6^\circ$. Figure 4E shows that, at higher slopes, ($\sim 24^\circ$) the reflections become independent of the roughness parameter and show the entire solar flux (I/F_{max} , see Method) distributed over Titan's hemisphere.

6. Results and Discussion

Based on our detailed analysis of the *Cassini* data coupled with roughness modeling and testing the feasibility of various possibilities, we propose that the BEF represents a broad specular reflection from a rough, wetted surface —presumably brought about by rainfall (T120 flyby, June 07, 2016), 2 Titan days after the observed cloud coverage (T119 flyby, May 06, 2016). A rough land surface could be smoothed by a recent surface wetting due to a summer-rainfall event. Earth analogs of wetted sidewalks (Figure 1E) and a mudflat [Clark *et al.*, 2010] in Utah's Bonneville Salt Flats (Figure 1F) broadly fit the BEF profile. The RADAR data show the area to be rough, variegated, and non-uniform (at scales of ~ 2 cm). A wet-sidewalk region is wet on a solid but rough substrate that at the right geometries produces broad specular reflections. These help us conclude that the

BEF would have properties similar to a wet sidewalk that arises from a recently wetted, solid and rough surface on the north pole of Titan.

7. Conclusion

The reduced and delayed cloud and storm activity at Titan's north pole as opposed to the model predicted activity suggests that our current understanding of Titan's changing seasons is incomplete. Our reported observations represent the first documented rainfall event on the north pole of Titan. We discover this rainfall based on a novel technique "wet-sidewalk effect" (broad specular reflections from a smoothed, original rough surface). Identifying the location and timing of the rainfall offers important data points for Titan GCMs, helping constrain why the northern summer cloud activity is delayed relative to predictions —especially because the north pole hosts most of Titan's lakes and seas [Hayes, 2016]. Apart from helping us understand Titan's weather and long term climate, a broad specular reflection observation like this helps us derive the roughness of the solid surfaces on Titan. Surface roughness of a region holds clues to its geologic evolution, nature of surface interactions and even has implications for landing site evaluations for future missions [Turtle *et al.*, 2017] to Titan.

Acknowledgments

The authors acknowledge support from the NASA/ESA Cassini Project. RD, JWB, CS and JS acknowledge support from NASA Cassini Data Analysis and Participating Scientists (CDAPS) grant NNX15AI77G. RD is grateful to Shannon MacKenzie, Deepak Dhingra, Johnathon Ahlers, Robert Chancia, Matthew Hedman and Subrata Chakraborty for manuscript suggestions, and would also like to thank the Cassini team

for all the years of profound work. The VIMS data are available at the PDS imaging node (<https://pds-imaging.jpl.nasa.gov/volumes/vims.html>). The RADAR data was obtained from (<http://pds-imaging.jpl.nasa.gov/volumes/radar.html>). Thanks to Dr. Jani Radebaugh and the other reviewer for their constructive comments and helpful suggestions on an earlier version of the manuscript.

References

- Ádámkóvics, M., M. H. Wong, C. Laver, and I. de Pater, Widespread Morning Drizzle on Titan, *Science*, *318*, 962–965, doi:10.1126/science.1146244, 2007.
- Barnes, J. W., et al., Global-scale surface spectral variations on Titan seen from Cassini/VIMS, *Icarus*, *186*, 242–258, doi:10.1016/j.icarus.2006.08.021, 2007.
- Barnes, J. W., et al., Organic sedimentary deposits in Titan’s dry lakebeds: Probable evaporite, *Icarus*, *216*, 136–140, doi:10.1016/j.icarus.2011.08.022, 2011.
- Barnes, J. W., et al., A Transmission Spectrum of Titan’s North Polar Atmosphere from a Specular Reflection of the Sun, *Astrophysical Journal*, *777*, 161, doi:10.1088/0004-637X/777/2/161, 2013a.
- Barnes, J. W., et al., Precipitation-Induced Surface Brightenings Seen on Titan by Cassini VIMS and ISS, *Planetary Science*, *2:1*, 2013b.
- Barnes, J. W., et al., Cassini/vims observes rough surfaces on titan’s punga mare in specular reflection, *Planetary Science*, *3(1)*, 3, doi:10.1186/s13535-014-0003-4, 2014.
- Brown, M. E., A. L. Smith, C. Chen, and M. Ádámkóvics, Discovery of Fog at the South Pole of Titan, *Astrophysical Journal*, *706*, L110–L113, doi:10.1088/0004-637X/706/1/L110, 2009.

- Brown, R. H., et al., The Cassini Visual And Infrared Mapping Spectrometer (Vims) Investigation, *Space Science Reviews*, *115*, 111–168, doi:10.1007/s11214-004-1453-x, 2004.
- Clark, R. N., et al., Detection and mapping of hydrocarbon deposits on Titan, *JGR Planets*, *115*(E14), 10,005, doi:10.1029/2009JE003369, 2010.
- Cottini, V., C. A. Nixon, D. E. Jennings, R. de Kok, N. A. Teanby, P. G. J. Irwin, and F. M. Flasar, Spatial and temporal variations in Titan's surface temperatures from Cassini CIRS observations, *PSS*, *60*, 62–71, doi:10.1016/j.pss.2011.03.015, 2012.
- Elachi, C., et al., Radar: The Cassini Titan Radar Mapper, *Space Science Reviews*, *115*, 71–110, doi:10.1007/s11214-004-1438-9, 2004.
- Griffith, C. A., et al., Characterization of Clouds in Titan's Tropical Atmosphere, *Astrophysical Journal*, *702*, L105–L109, doi:10.1088/0004-637X/702/2/L105, 2009.
- Hayes, A. G., The lakes and seas of titan, *Annual Review of Earth and Planetary Sciences*, *44*(1), 57–83, doi:10.1146/annurev-earth-060115-012247, 2016.
- Hofgartner, J., et al., Transient features in a titan sea, *Nature geoscience*, 2014.
- Jennings, D. E., et al., Composite infrared spectrometer (cirs) on cassini: publisher's note, *Appl. Opt.*, *56*(21), 5897–5897, doi:10.1364/AO.56.005897, 2017.
- MacKenzie, S. M., et al., Evidence of titan's climate history from evaporite distribution, *Icarus*, *243*(0), 191 – 207, doi:http://dx.doi.org/10.1016/j.icarus.2014.08.022, 2014.
- Malaska, M. J., R. Hodyss, J. I. Lunine, A. G. Hayes, J. D. Hofgartner, G. Hollyday, and R. D. Lorenz, Laboratory measurements of nitrogen dissolution in Titan lake fluids, *Icarus*, *289*, 94–105, doi:10.1016/j.icarus.2017.01.033, 2017.

McCord, T. B., et al., Composition of Titan's surface from Cassini VIMS, *PSS*, *54*, 1524–1539, 2006.

Porco, C. C., et al., Cassini Imaging Science: Instrument Characteristics And Anticipated Scientific Investigations At Saturn, *Space Science Reviews*, *115*, 363–497, 2004.

Porco, C. C., et al., Imaging of Titan from the Cassini spacecraft, *Nature*, *434*, 159–168, 2005.

Rodriguez, S., et al., Cassini/VIMS hyperspectral observations of the HUYGENS landing site on Titan, *PSS*, *54*, 1510–1523, 2006.

Rodriguez, S., et al., Global circulation as the main source of cloud activity on Titan, *Nature*, *459*, 678–682, doi:10.1038/nature08014, 2009.

Rodriguez, S., et al., Titan's cloud seasonal activity from winter to spring with Cassini/VIMS, *Icarus*, *216*, 89–110, doi:10.1016/j.icarus.2011.07.031, 2011.

Roe, H. G., I. de Pater, B. A. Macintosh, and C. P. McKay, Titan's Clouds from Gemini and Keck Adaptive Optics Imaging, *Astrophysical Journal*, *581*, 1399–1406, 2002.

Schaller, E. L., M. E. Brown, H. G. Roe, and A. H. Bouchez, A large cloud outburst at titan's south pole, *Icarus*, *182*(1), 224–229, 2006.

Schaller, E. L., M. E. Brown, H. G. Roe, A. H. Bouchez, and C. A. Trujillo, Dissipation of titan's south polar clouds, *Icarus*, *184*(2), 517–523, 2006.

Schneider, T., S. D. B. Graves, E. L. Schaller, and M. E. Brown, Polar methane accumulation and rainstorms on Titan from simulations of the methane cycle, *Nature*, *481*, 58–61, doi:10.1038/nature10666, 2012.

- Soderblom, J. M., et al., Modeling specular reflections from hydrocarbon lakes on Titan, *Icarus*, *220*, 744–751, doi:10.1016/j.icarus.2012.05.030, 2012.
- Sotin, C., et al., Observations of Titan's Northern lakes at 5 μm : Implications for the organic cycle and geology, *Icarus*, *221*, 768–786, doi:10.1016/j.icarus.2012.08.017, 2012.
- Stephan, K., et al., Specular reflection on Titan: Liquids in Kraken Mare, *Geophysical Research Letter*, *37*, L7104, doi:10.1029/2009GL042312, 2010.
- Tokano, T., and R. D. Lorenz, GCM simulation of balloon trajectories on Titan, *PSS*, *54*, 685–694, doi:10.1016/j.pss.2006.04.001, 2006.
- Tokano, T., C. P. McKay, F. M. Neubauer, S. K. Atreya, F. Ferri, M. Fulchignoni, and H. B. Niemann, Methane drizzle on Titan, *Nature*, *442*, 432–435, doi:10.1038/nature04948, 2006.
- Turtle, E. P., J. E. Perry, A. S. McEwen, A. D. Del Genio, J. Barbara, R. A. West, D. D. Dawson, and C. C. Porco, Cassini imaging of Titan's high-latitude lakes, clouds, and south-polar surface changes, *Geophysical Research Letters*, *36*, L2204, doi:10.1029/2008GL036186, 2009.
- Turtle, E. P., et al., Rapid and Extensive Surface Changes Near Titan's Equator: Evidence of April Showers, *Science*, *331*, 1414–1417, doi:10.1126/science.1201063, 2011.
- Turtle, E. P., et al., Dragonfly: Exploring Titan's Prebiotic Organic Chemistry and Habitability, in *Lunar and Planetary Science Conference, Lunar and Planetary Science Conference*, vol. 48, p. 1958, 2017.

Table 1. Details of T106, T120 and T121 flybys, which cover the location of BEF over three time stamps. It helps us determine the existence of this feature over time, and we conclude that it is an ephemeral feature. The inferred specular points of the flybys are tabulated. We also show the distance of T106 and T121 specular points from the BEF. Sp. Pt. means Specular Point. The fact that the distance of inferred specular point

Flyby Info	Cube No	Pixel Sampling (pixel/degree)	Phase (°)	Incidence (°)	Emission (°)	Inferred Sp. Pt.	Dist. from BEF (km)	Spatial Sampling (lat x long) (km/pixel)
T106, 24 Oct. 2014	CM1792831393	2.2	120	50	73	71°N, 65°W	847	100 x 90
T120, 07 June 2016	CM11844022476	2.08	116	52	65	82°N, 78°W	632	55 x 70
T121, 25 July 2016	CM11848148220	3.1	113	65	55	83°N, 6°W	470	55 x 35




Figure 1. VIMS wet-sidewalk color composite (R:5 μ m, G: 2.7 μ m, B:2 μ m) of Titan's north polar region showing the region corresponding to the BEF in the flybys A) T106, 24 October, 2014 B) T120, 07 June, 2016, and C) T121, 25 July, 2016. The yellow box in 1C shows the disappearance of the Bright Ephemeral Feature (BEF) three Titan days later (See Table 1). D) Zoomed- in view of the north polar region from the T120 flyby annotated to identify the features. The BEF is marked by magenta arrows in the figure. The black dot marks the north pole. E) Analog for the T120 BEF on a cloudy day after rainfall. The specular reflection at the boundary of a puddle can be seen as a bright speck while the road's wetted surface is reflecting broadly at the right geometries. F) Analog for the T120 BEF as mudflat in Utah's Bonneville salt flats




Figure 2. The figure is in a different orientation than Figure 1. Kraken Mare is in the bottom right as opposed to the bottom left in the previous figures. A) RADAR map of the north polar region of Titan in Polar Stereographic projection. The yellow box indicates the region corresponding to the BEF illustrating that the region is mostly solid land. The non-data regions are white. B) VIMS infrared color composite (R:5 μm , G: 2.7 μm , B:1.3 μm) representing the yellow-boxed region as in the RADAR map of the north pole. C) VIMS infrared color composite (3B) (R:5 μm , G: 2.7 μm , B:1.3 μm) overlain on the north polar map of RADAR (3A) showing the BEF and specular point. The red arrow indicates the specular reflection point (82.7°N, 78.6°W) near Xolotlan Lacus. The magenta arrows indicate the BEF we observe in T120.




Figure 3. A) The colored boxes correspond to the different regions in the T120 observations: dark blue (clouds), green (land), light blue (sea), red (specular reflection), magenta (BEF), and yellow (evaporites). B) The spectra corresponding to the regions in similar colors as the boxes in A. Grey solid boxes show spectral regions affected by methane absorption and thus blocked by Titan’s atmosphere. C) Shows the spectra for the location of the BEF. The slightly higher reflectance at shorter wavelengths of the T106 observation (cyan) is because of the higher emission angles. The magenta spectrum corresponds to the BEF during the T120 June 2016 flyby which is distinctly brighter than other spectra at 5 μ m. The yellow spectrum indicates reflectances for the same region three Titan days later during the T121 July 2016 flyby.




Figure 4. A) Modeled reflectance (I/F) variation with surface roughness for the broad specular reflection (BEF) observed during the T120 flyby. The modeled (magenta) reflectance peaks at roughness (σ) values of $\sim 14^{\circ}$ indicating that the BEF’s surface roughness could be of that order. B) T120 *Cassini* image cube CM1866022476 where the BEF is observed. C) Model-generated synthetic images for the T120 observation geometry with a surface roughness parameter of 2° reveals only the specular reflection. One thing to note is the scale in the synthetic images: as we increase the roughness or specular deviation angle — σ —the specular reflection spreads over a larger number of pixels, each pixel receiving lesser solar flux for higher σ and hence the scale changes. D) σ of 12° covers the extent of BEF. E) σ of 24° distributes the entire solar flux onto the Titan hemisphere.